

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

**(19) World Intellectual Property Organization
International Bureau**



A standard linear barcode is located at the bottom of the page, spanning most of the width.

**(43) International Publication Date
7 April 2005 (07.04.2005)**

PCT

**(10) International Publication Number
WO 2005/031365 A3**

(51) International Patent Classification⁷: G01N 33/68,
C07K 14/475, G01N 33/543

GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW

(21) International Application Number: PCT/IB2004/003606

Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(22) International Filing Date: 29 September 2004 (29.09.2004)

kind

(25) Filing Language: English

GM,

(26) Publication Language: English

Europ

(30) Priority Data: 60/507,192 29 September 2003 (29.09.2003) US

Published:

— with international search report

(88) Date of publication of the international search report:
7 July 2005

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2005/031365 A3

(54) Title: RAPID DETERMINATION OF HEPATOCYTE GROWTH FACTOR (HGF) IN THE BODY FLUIDS

(57) Abstract: A method for diagnosing an inflammatory disorder in an individual comprises contacting a body fluid sample from the individual with an immunoassay device to determine relative levels of different forms of hepatocyte growth factor (HGF) in the sample, and correlating the determined levels to an inflammatory disorder. An immunoassay device for diagnosing an inflammatory disorder in an individual comprises at least two test areas having immobilized respectively therein reagent for binding different forms of hepatocyte growth factor (HGF).